

CONTENTS

Special issue: A collection of papers on liquid chromatography

Foreword	
K. B. Hicks (Philadelphia, PA, U.S.A.)	vii
High-performance anion-exchange chromatography of sugars and sugar alcohols on quaternary ammonium resins under alkaline conditions	
T. J. Paskach, P. J. Reilly (Ames, IA, U.S.A.), H.-P. Lieker, and K. Thielecke (Braunschweig, Germany)	1
H.p.l.c. of 4-nitrophenyl- α -malto-oligosaccharides	
C. Niemann, W. Saenger, R. Nuck (Berlin, F.R.G.), and B. Pfannemüller (Freiburg, Germany)	15
High performance reversed-phase chromatographic mapping of 2-pyridylamino derivatives of xyloglucan oligosaccharides	
Z. El Rassi, D. Tedford, J. An, and A. Mort (Stillwater, OK, U.S.A.)	25
Selectivity control in the anion-exchange chromatographic determination of saccharides in dairy products using pulsed amperometric detection	
J. van Riel and C. Olieman (Ede, The Netherlands)	39
High-performance size-exclusion chromatography and ELISA detection of extracellular polysaccharides from <i>Mucorales</i>	
G. A. de Ruiter, A. W. van der Lugt, A. G. J. Voragen and F. M. Rombouts (Wageningen, The Netherlands) and S. H. W., Notermans (Bilthoven, The Netherlands)	47
An h.p.l.c. method for determining chain-length distribution in some glycogens	
N. W. H. Cheetham, N. Hansawek and P. Saecou (Kensington, Australia)	59
High-performance liquid chromatography of mono- and oligo-saccharides on a graphitized carbon column	
K. Koizumi, Y. Okada, and M. Fukuda (Nishinomiya, Japan)	67
Isolation of oligogalacturonic acids in gram quantities by preparative h.p.l.c.	
A. Hotchkiss, Jr., K. B. Hicks, L. W. Doner and P. L. Irwin (Philadelphia, U.S.A.)	81
Intrinsic viscosity and molecular weight of pectin components	
M. L. Fishman, D. T. Gillispie, and S. M. Sondey (Philadelphia, U.S.A.) and Y. S. El-Atawy (Giza, Egypt)	91
Analysis of oligouronates by reversed-phase ion-pair h.p.l.c.: Role of the mobile phase	
A. Heyraud and C. Leonard (Grenoble, France)	105
Determination of furosine in biomedical samples employing an improved hydrolysis and high-performance liquid chromatographic technique	
W. T. Cefalu, A. Bell-Farrow, Z. Q. Wang, and S. Ralapati (Winston-Salem, NC, U.S.A.)	117
Characterization of five isomers of branched cyclomaltoheptaose (β CD) having degree of polymerization (d.p.) = 9: Reinvestigation of three positional isomers of diglucosyl- β CD	
K. Koizumi, T. Tanimoto, Y. Okada, N. Nakanishi, N. Kato (Nishinomiya, Japan), Y. Takagi and H. Hashimoto (Yokohama, Japan)	127

Kinetic analysis of pectate lyases by high-performance liquid chromatography J. F. Preston III and J. D. Rice (Gainesville, FL, U.S.A.)	137
Kinetic comparisons of trimer-generating pectate and alginate lyases by reversed-phase ion-pair liquid chromatography J. F. Preston III, J. D. Rice, M. C. Chow, and B. J. Brown (Gainesville, FL, U.S.A.) . . .	147
Optimization of waveforms for pulsed amperometric detection (p.a.d.) of carbohydrates following separation by liquid chromatography W. R. LaCourse and D. C. Johnson (Ames, IA, U.S.A.)	159
High-performance anion-exchange chromatography with pulsed amperometric detection of linear and branched glucose oligosaccharides R. N. Ammeraal, G. A. Delgado, F. L. Tenbarge, and R. B. Friedman (Hammond, IN, U.S.A.)	179
Capillary zone electrophoresis of reducing mono- and oligo-saccharides as the borate complexes of their 3-methyl-1-phenyl-2-pyrazolin-5-one derivatives S. Honda, S. Suzuki, A. Nose, K. Yamamoto, and K. Kakehi (Higashiosaka, Japan) . . .	193
<i>Notes</i>	
Chromatography of 1-deoxy-1-nitroalditols on a cation-exchange resin in the lanthanum form M. Petrušová, L. Petruš, and E. Lattová (Bratislava, Czechoslovakia)	199
Analyses of α -linked disaccharides of D-glucose by high-performance liquid chromatography M. Shiota (Tokyo, Japan) and S. Kobayashi (Ibaraki, Japan)	203
Separation of high-mannose isomers from yeast and mammalian sources using high-pH anion-exchange chromatography R. R. Townsend (San Francisco, CA, U.S.A.), P. H. Atkinson (Bronx, NY, U.S.A.), and R. B. Trimble (Albany, NY, U.S.A.)	211
Problems encountered during the extraction, purification, and chromatography of pectic fragments, and some solutions to them A. J. Mort, B. M. Moerschbacher, M. L. Pierce and N. O. Maness (Stillwater, OK, U.S.A.)	219
<i>End of special issue papers</i>	
Molecular motion in glassy water-malto-oligosaccharide (maltodextrin) mixtures as studied by conventional and saturation-transfer spin-probe e.s.r. spectroscopy M. J. G. W. Roozen, M. A. Hemminga and P. Walstra (Wageningen, Netherland)	229
The conformation of 1,6-anhydrolactose and its hexa-acetate in solution A. Rivera-Sagredo and J. Jiménez-Barbero (Madrid, Spain)	239
A molecular modelling study of the distortion of α -cyclodextrin (cyclomaltohexaose) in complexes with guest molecules S. P. van Helden, M. J. van Drooge, A. J. Claessens, A. C. A. Jansen, and L. H. M. Janssen (Utrecht, The Netherlands)	251
Synthesis of disaccharides with (1 \rightarrow 4)- β glycosidic linkages related to the 4- and 6-sulfates and the 4,6-disulfates of chondroitin M. Zšiška (Oldenburg, Germany) and B. Meyer (Athens, GA, U.S.A.)	261
Synthesis of β -D-GlcA-(1 \rightarrow 3)- β -D-Gal disaccharides with 4- and 6-sulfate groups and 4,6-disulfate groups M. Zšiška (Oldenburg, Germany) and B. Meyer (Athens, GA, U.S.A.)	279

Structure of an acidic glycan present in the lipopolysaccharide extract from the reference strain for <i>Serratia marcescens</i> serogroup O18 D. Oxley and S. G. Wilkinson (Hull, Gt. Britain)	293
Structural studies of the <i>Vibrio cholerae</i> O:3 O-antigen polysaccharide T. A. Chowdhury, P.-E. Jansson, B. Lindberg, J. Lindberg, B. Gustafsson, and T. Holme (Stockholm, Sweden)	303
Distinguishing mammalian sialidases by inhibition kinetics with novel derivatives of 5-acetamido-2,6-anhydro-3,5-dideoxy-D-glycero-D-galacto-non-2-enonic acid, an unsat- urated derivative of <i>N</i> -acetylneuraminic acid T. G. Warner (Memphis, TN, U.S.A.), A. Louie (South San Francisco, CA, U.S.A.), M. Potier (Montreal, Canada), A. Ribeiro (Durham, NC, U.S.A.)	315
Structural analysis of the heptose/hexose region of the lipopolysaccharide from <i>Escherichia coli</i> K-12 strain W3100 O. Holst, U. Zähringer, H. Brade (Borstel, Germany), and A. Zamojski (Warsaw, Poland)	323
<i>Notes</i>	
The synthesis and characterisation of 6- <i>O</i> -L-glycero-D-manno-heptopyranosyl-D-glucopyranose Z. Pakulski, A. Zamojski (Warsaw, Poland), O. Holst, and U. Zähringer (Borstel, Germany)	337
Configurational assignment of <i>N</i> -acetylneuraminic acid and analogues <i>via</i> the vicinal C,H coupling constants S. Prytulla, J. Lauterwein, M. Klessinger, and J. Thiem (Münster, Germany)	345
Determination of the activation energy and rate constants for the chair-to-chair interconversion of <i>cis</i> -inositol by n.m.r. spectroscopy K. Dill and L. Huang (Clemson, SC, USA)	351
Complexing of sugars and sugar alcohols with metal ions: a comparative study by ion-exchange chromatography M. M. Hämäläinen and H. Lönnberg (Turku, Finland)	357
<i>Preliminary communication</i>	
Iodonium ion-assisted glycosylation of alkyl (aryl) 1-thioglycosides: regulation of stereoselectivity and reactivity H. M. Zuurmond, S. C. van der Laan, G. A. van der Marel, and H. van Boom (Leiden, The Netherlands)	c1
<i>Author index</i>	c5
<i>Subject index</i>	c7
<i>Contents (Vol. 215)</i>	c11

